

12-23-2013

Correlation of *Aggregatibacter actinomycetemcomitans* Detection with Clinical/Immunoinflammatory Profile of Localized Aggressive Periodontitis Using a 16S rRNA Microarray Method: A Cross-Sectional Study

Patricia F. Gonçalves

Vanja Klepac-Ceraj
vklepacc@wellesley.edu

Hong Huang

Bruce J. Paster

Ikramuddin Aukhil

See next page for additional authors

Follow this and additional works at: <http://repository.wellesley.edu/scholarship>

Version: Publisher's version

Recommended Citation

Gonçalves PF, Klepac-Ceraj V, Huang H, Paster BJ, Aukhil I, et al. (2013) Correlation of *Aggregatibacter actinomycetemcomitans* Detection with Clinical/Immunoinflammatory Profile of Localized Aggressive Periodontitis Using a 16S rRNA Microarray Method: A Cross-Sectional Study. PLoS ONE 8(12): e85066. doi:10.1371/journal.pone.0085066

This Article is brought to you for free and open access by Wellesley College Digital Scholarship and Archive. It has been accepted for inclusion in Faculty Research and Scholarship by an authorized administrator of Wellesley College Digital Scholarship and Archive. For more information, please contact ir@wellesley.edu.

Authors

Patricia F. Gonçalves, Vanja Klepac-Ceraj, Hong Huang, Bruce J. Paster, Ikramuddin Aukhil, Shannon M. Wallet, and Luciana M. Shaddox

Correlation of *Aggregatibacter actinomycetemcomitans* Detection with Clinical/Immunoinflammatory Profile of Localized Aggressive Periodontitis Using a 16S rRNA Microarray Method: A Cross-Sectional Study

Patricia F. Gonçalves^{1,2}, Vanja Klepac-Ceraj^{3,4}, Hong Huang², Bruce J. Paster^{3,5}, Ikramuddin Aukhil², Shannon M. Wallet², Luciana M. Shaddox^{2*}

1 Department of Dentistry, Federal University of Jequitinhonha and Mucuri Valleys, Diamantina, Minas Gerais, Brazil, **2** Department of Periodontology, University of Florida College of Dentistry, Gainesville, Florida, United States of America, **3** Department of Microbial Ecology and Pathogenesis, The Forsyth Institute, Cambridge, Massachusetts, United States of America, **4** Department of Biological Sciences, Wellesley College, Wellesley, Massachusetts, United States of America, **5** Department of Oral Medicine, Infection and Immunity, Harvard School of Dental Medicine, Boston, Massachusetts, United States of America

Abstract

Objective: The objective of this study was to determine whether the detection of *Aggregatibacter actinomycetemcomitans* (Aa) correlates with the clinical and immunoinflammatory profile of Localized Aggressive Periodontitis (LAP), as determined by 16S rRNA gene-based microarray.

Subjects and Methods: Subgingival plaque samples from the deepest diseased site of 30 LAP patients [PD \geq 5 mm, BoP and bone loss] were analyzed by 16S rRNA gene-based microarrays. Gingival crevicular fluid (GCF) samples were analyzed for 14 cyto/chemokines. Peripheral blood was obtained and stimulated in vitro with *P.gingivalis* and *E.coli* to evaluate inflammatory response profiles. Plasma lipopolysaccharide (LPS) levels were also measured.

Results: Aa was detected in 56% of LAP patients and was shown to be an indicator for different bacterial community structures ($p < 0.01$). Elevated levels of pro-inflammatory cyto/chemokines were detected in LPS-stimulated blood samples in both Aa-detected and Aa-non-detected groups ($p > 0.05$). Clinical parameters and serum LPS levels were similar between groups. However, Aa-non-detected GCF contained higher concentration of IL-8 than Aa-detected sites ($p < 0.05$). TNF α and IL1 β were elevated upon *E.coli* LPS stimulation of peripheral blood cells derived from patients with Aa-detected sites.

Conclusions: Our findings demonstrate that the detection of Aa in LAP affected sites, did not correlate with clinical severity of the disease at the time of sampling in this cross-sectional study, although it did associate with lower local levels of IL-8, a different subgingival bacterial profile and elevated LPS-induced levels of TNF α and IL1 β .

Citation: Gonçalves PF, Klepac-Ceraj V, Huang H, Paster BJ, Aukhil I, et al. (2013) Correlation of *Aggregatibacter actinomycetemcomitans* Detection with Clinical/Immunoinflammatory Profile of Localized Aggressive Periodontitis Using a 16S rRNA Microarray Method: A Cross-Sectional Study. PLoS ONE 8(12): e85066. doi:10.1371/journal.pone.0085066

Editor: Paul J Planet, Columbia University, United States of America

Received: July 8, 2013; **Accepted:** November 22, 2013; **Published:** December 23, 2013

Copyright: © 2013 Gonçalves et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was supported by NIH/NIDCR R01DE019456 (www.nidcr.nih.gov). PF Gonçalves postdoctoral fellowship was supported by the National Council of Technological and Scientific Development, CNPq, Brazil (www.cnpq.br). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

* E-mail: lshaddox@dental.ufl.edu

Introduction

Aggregatibacter actinomycetemcomitans (Aa) is a Gram-negative bacterium that colonizes the oral cavity of one third or more of the population aged up to 18 years [1]. It has been highly implicated as the causative agent of aggressive periodontitis (AgP), a group of less frequent, often severe, rapidly progressive forms of periodontitis, with a more localized

presentation of the disease (LAP) [2–4]. LAP is also characterized by an early age of onset, tendency for familial involvement, and bone loss affecting specifically molar and incisor teeth, [5]. Importantly, recent papers have pointed out the importance Aa, a particularly highly leukotoxic clone (JP2), in the initiation of attachment loss in young individuals [4,6,7].

Our group has recently demonstrated a hyper-inflammatory trait in response to bacterial endotoxin in an LAP cohort of

African-Americans, which results in the over-expression of pro-inflammatory cyto/chemokines [8]. We also observed an exacerbated local inflammatory response, along with high systemic levels of bacterial lipopolysaccharides (LPS) [9]. Additionally, we have reported Aa to be highly associated with LAP in this cohort [10]. However, we could not detect Aa in every patient with LAP in our cohort, which is also consistent to previously reported literature [11]. Little is known regarding the impact of Aa's presence on LAP disease presentation (clinical parameters and bacterial colonization) or the accompanying inflammatory response (regulation and local presentation).

Thus, the objective of this study was to evaluate our LAP cohort in relation to the presence of Aa and determine whether the detection of this species in LAP affected sites correlate with clinical presentation of disease, local cyto/chemokines, plasma LPS levels and the level of hyper-responsiveness of these individuals.

Materials and Methods

Ethics statement

This study was conducted in full conformity with the principles set forth in The Belmont Report: Ethical Principles and Guidelines for the Protection of Human Subjects of Research, as drafted by the US National Commission for the Protection of Human Subjects of Biomedical and Behavioral Research (April 18, 1979) and codified in 45 CFR Part 46 and/or the ICH E6; 62 Federal Regulations 25691 (1997). This study was approved by the University of Florida Institutional Review Board.

Subject demographics

Subjects were selected from a cohort recruited from Leon County Health Department, Tallahassee, Florida, from February 2007 to August 2010. This study is part of a larger clinical trial (Clinical trial registration: #NCT01330719 at www.clinicaltrials.gov). A comprehensive microbiological profile as well as local and LPS-induced inflammatory response from initial subjects enrolled in this study has been reported in our previous publications [8-10]. Both historical data and newly generated data were part of the present analysis. Specifically, all subjects enrolled in the clinical trial that had their baseline microbiological analysis completed were included in the present report [10], and new data on local and systemic inflammatory markers for some of these patients were run to fulfill the present investigation (see flowchart, Figure 1). Eight of the participants had siblings within the cohort, who happen to be part of the random pool of participants on which microbiological analysis had been completed. All subjects and parents were informed about the study protocol and signed an informed consent previously approved by the University of Florida Institutional Review Board. Inclusion criteria: subjects aged 5-21 years old, African-American, diagnosed with localized aggressive periodontitis, defined by presence of at least 2 sites, on first molar and/or incisor, with pocket depth >4mm with bleeding on probing, presence of at least 2mm clinical attachment loss and radiographic bone loss [5,12]. Exclusion criteria: subjects diagnosed with any systemic




Figure 1. Schematic flowchart of study design.

doi: 10.1371/journal.pone.0085066.g001

diseases or conditions that could influence the progression and/or clinical characteristics of periodontal disease (e.g., diabetes or blood disorders); taken antibiotics within the last 3 months or any medications that could influence the characteristics of the disease (e.g., phenytoin, cyclosporine); have received periodontal treatment within the previous 6 months; smokers; and pregnant/lactating women.

Clinical measurements

The following periodontal clinical parameters were taken by one calibrated examiner at the baseline visit (prior to treatment): Pocket Depth (PD), Bleeding on probing (BoP); Gingival margin position (GM), Clinical attachment level (CAL); Plaque Index (PI). All measurements were performed using a UNC-15 periodontal probe at six sites per tooth and recorded using a computer software program (Florida Probe, Gainesville, FL, USA). Periapical and interproximal radiographs were taken on all patients at the baseline visit for diagnosis purposes only.

Gingival Crevicular Fluid (GCF) sampling

Following clinical examination, and on the same day, samples were collected from the most severe diseased site (PD \geq 5mm with BoP and CAL \geq 2 mm and radiographic bone loss). Sites for GCF collection were isolated with cotton rolls, gently air-dried and supragingival plaque removed. A collection strip (PerioPaper GCF collection strips, Oraflow Inc, Plainview, NY) was inserted in the sites 1-2 mm into the sulcus for ~10 sec, as described previously [9]. Volume of GCF was measured using a Periotron[®]8000, (Oraflow, Inc.). GCF samples were frozen at -70°C until analysis of soluble mediators was performed.

Collection of bacterial sub-gingival biofilm

Subgingival plaque samples were collected from the same diseased site where GCF collection was made, as described above, also prior to treatment. The area of collection was isolated with cotton rolls and supragingival plaque was carefully removed. A sterile endodontic paper point was used for the

collection, placed in an eppendorf tube and stored at -70°C until processed.

Microbial microarray analysis

DNA was isolated from plaque samples as previously described [10]. After purification, DNA concentration and quality was determined using the Nanodrop (ND-1000 Spectrophotometer, Nanodrop Technologies INC, Wilmington, DE, USA). Samples were analyzed at the Forsyth Institute using the Human Oral Microbe Identification Microarrays (HOMIM) as previously described [10,13]. Briefly, 16S rRNA genes were PCR amplified from DNA extracts using 16S rRNA universal forward and reverse primers and labeled via incorporation of Cy3-dCTP in a second nested PCR. HOMIM uses 16S rRNA-based, reverse-capture oligonucleotide probes (typically 18 to 20 bases), which are printed on aldehyde-coated glass slides and probed with labeled PCR products described above which are hybridized in duplicate. The microarray slides are scanned using an Axon 4000B scanner and crude data is extracted using GenePix Pro software (Molecular Devices, Sunnyvale, CA). Microbial profiles were generated from image files of scanned arrays using a HOMIM online analysis tool (<http://bioinformatics.forsyth.org/homim/>). Detection of a particular taxon was determined by presence of a fluorescent spot for that unique probe. A mean intensity for each taxon was calculated from hybridization spots of the same probe and signals were normalized by comparing individual signal intensities to the average of signals from universal probes and calculated as described previously [13]. Any original signal that was less than two times the background value was reset to 1 and was assigned to the signal level 0. Signals greater than 1 were categorized into scores from 1 to 5, corresponding to ranked signal levels. Presence of Aa was determined if a site showed value of 1 or above. Absence of Aa was determined if a site showed a value of '0'

In vitro LPS stimulation

Peripheral blood was collected in heparinized tubes, diluted 1:4 in RPMI1640 (Invitrogen, Carlsbad, CA, USA) and stimulated with 1 $\mu\text{g}/\text{mL}$ of ultrapure *E.coli* (Ec) LPS or *P.gingivalis* (Pg) LPS (InvivoGen, San Diego, CA, USA), or left untreated, for 24hrs [8]. After which the resulting supernatants were collected and stored at -70°C until analysis of soluble mediators was performed.

Analysis of soluble mediators

Multiplex assays (Milliplex[®] 14-plex cytochemokine detection kits, Millipore, St. Charles, MO, USA) were used to detect and quantify 14 cyto/chemokines (Eotaxin, TNF α , IFN γ , IL1 β , IL2, IL6, IL7, IL8, IL10, IL12p40, MCP-1, G-CSF, GM-CSF and MIP1 α) in the crevicular fluid and LPS-stimulated culture supernatants according to the manufacturer's protocol. Data was acquired on the Luminex[®]200[™] (Millipore, St. Charles, MO) and analyzed with Milliplex Software (Viagene Tech, Carlisle, MA, USA), a standard curve and 5-parameter logistics. Data is reported as pg/ml. GCF data is normalization to total protein content: normalized pg/ml = [pg/ml x protein

content corrective ratio], where corrective ratio = [lowest protein concentration/protein concentration of sample of interest].

Plasma LPS levels

Peripheral blood was collected in heparinized tubes, after which plasma was separated from red blood cells by centrifugation ($\sim 300 \times g$ for 15 min) and stored at -70°C . LPS levels were detected and quantified by using a chromogenic assay (Endpoint Chromogenic LAL Assay, Lonza, Basel, Switzerland). Endotoxin units/ml were calculated using a standard curve and best fit linear trend line, as previously described [9].

Statistical analysis

Means were calculated and t-test was performed to compare patients presenting *Aa-detected* sites versus patients presenting *Aa-non-detected* sites for clinical and immunoinflammatory parameters. Mann-Whitney Rank Sum Test was applied when data was not normally distributed. ANOVA analyses were used for comparisons within groups for LPS stimulation using different ligands. This study was powered at $>80\%$ to detect a 10% difference between the two groups in cyto/chemokine values [values based on previous publication data [8,9]] and a PD difference in affected sites of at least 1 mm with standard deviation of 0.9 mm. SigmaPlot version 11.0 software (Systat Software Inc, Chicago, IL) was used for statistical analysis, considering a 5% significance level. We considered cyto/chemokines to be independent variables and thus, no statistical corrections were done here [14]. Canonical correspondence analysis (CCA), nonmetric multidimensional scaling (NMDS), and ANOVA tests were used on HOMIM hybridization scores to test for significant effects of environmental variables on microbial community structures within sites with or without the detection of Aa, using the Vegan package for the R statistical programming environment [15]. For generating CCA plots, Aa was excluded from the list of taxa. To account for the multiple hypotheses, we adjusted p-value using Benjamini/Hochberg correction [16].

Results

Clinical and microbiologic characterization

HOMIM resulted in 189 probes identifying at least 137 different bacterial taxa in diseased sites of LAP (for a more comprehensive microbial data description, including healthy controls, see [10]). Among these, two taxa of Aa were identified: *Aa_ot531_AA84* and *Aa_ot531_P02*. Sites presenting any signal for Aa (value 1 or greater) were considered *Aa-detected* (n=17) and those that did not present any signal for Aa (value 0) were considered *Aa-non-detected* (n=13). HOMIM analysis' mean relative signal intensity of Aa in the *Aa-detected* sites was 3.81 ± 1.02 , in a 0 to 5 signal scale. Interestingly, clinical parameters and demographics were similar between individuals with *Aa-detected* or *Aa-non-detected* sites, as determined by HOMIM (Table 1).

In NMDS and CCA plots (Figure 2), samples with similar community composition cluster close together, while samples

Table 1. Clinical and demographic parameters for Aa+(positive) and Aa-(negative) groups.

	Age	M/F	%PD>4mm	%BoP	%Plaque	Mean PD sites/all(mm)	Mean CAL(mm)	PD site (mm)
Aa+	13.19±3.94	8m/9f	12.76±7.81	19.41±10.05	43.64±25.88	5.02±0.71 / 2.27±0.40	3.94±1.53	5.94±1.19
Aa-	15.85±2.67	6m/7f	15.69±11.60	32.84±49.08	45.92±26.85	4.89±0.70 / 2.45±0.44	3.29±1.43	6.61±1.55
P value	0.371 [†]	0.891 [†]	0.416 [*]	0.834 [†]	0.986 [†]	0.608 [*] /0.258 [*]	0.243 [†]	0.220 [†]

Values are given by Means ± Standard deviation. M=male; F=female; Mean PD sites= Mean pocket depth of sites with PD>4mm; Mean PD all= mean PD of all sites; CAL=clinical attachment level of affected sites; BoP=bleeding on probing; PD site = Pocket depth from site sampled for microbiological analysis. *t-test; [†]Mann-Whitney Rank Sum Test.

doi: 10.1371/journal.pone.0085066.t001

with different community compositions are farther apart on the plots; preserving the rank order of the original similarities [17]. Both NMDS and CCA analysis suggest that the detection of Aa was associated with specific bacterial community structures (of all variables tested, only the presence of Aa was statistically significant, permutation test under the reduced model with $p<0.01$). PD was not statistically significant relative to Aa detection ($p=0.24$) and therefore was not included in CCA. The left side of the CCA ordination plot is predominantly occupied by diseased sites containing Aa, whereas on the right side are predominantly diseased sites in which Aa was not detected. We observed that *Granulicatella elegans* ($p=0.01$) and *Prevotella loeschii* ($p=0.02$) were frequently found in samples co-existing when Aa was present. In contrast, *Porphyromonas gingivalis* ($p=0.02$) and *Parvimonas micra* ($p=0.03$) were more likely to be found in Aa-non-detected communities. These significances disappear when corrected for multiple comparisons.

Local inflammatory profile and systemic LPS levels

As previously published, high levels of several inflammatory mediators were detected in the GCF of diseased sites of LAP participants [9]. Eotaxin, G-CSF and IL7 were not detected in high enough levels and thus were not included in the association analysis. In general, no differences in the GCF mediator profiles were detected between Aa-detected and Aa-non-detected sites (Figure 3). However, IL8 was found at higher concentrations in Aa-non-detected sites than those observed in Aa-detected sites ($p=0.011$). After controlling values for GCF volume at each diseased site of collection, the same trends were observed, and p-values slightly decreased, but did not reach statistical significance. Systemic LPS levels were similar in Aa-non-detected and Aa-detected groups (Mean values $0.61±0.15$ and $0.68±0.36$, respectively; $p=0.773$).

LPS-induced hyper-responsive phenotype

The LPS responsiveness of individuals with Aa-non-detected and Aa-detected groups was compared. There was no significant differences between the levels of cyto/chemokines in unstimulated cultures of Aa-detected and Aa-non-detected individuals. Pg and Ec LPS-stimulation of whole blood cell preparation resulted in elevated levels of multiple pro-inflammatory cyto/chemokines when compared with unstimulated cultures ($p<0.05$). There were few significant differences in the levels of cyto/chemokines induced when

comparing cohorts which had Aa-detected or Aa-non-detected sites. Specifically, Ec LPS stimulation elicited more robust levels of TNF α and IL1 β from peripheral blood of individuals with Aa-detected sites than those with Aa-non-detected sites ($p<0.05$) (Figure 4). In addition, GM-CSF was more highly induced by Pg LPS-stimulation in individuals with Aa-non-detected sites, compared to those with Aa-detected sites, although the difference did not reach significance (Figure 4).

Discussion

It has been previously suggested that an individual organism can play a strong role in community restructuring, and in turn impact the ecological processes of the system [18,19]. This change in community structure can also have an effect on local immune responses and thus the course of periodontal disease progression. Among periodontal pathogens, Aa has received particular attention and has been regarded as a key factor in the etiology of LAP. This view was based primarily on classic association studies linking Aa to disease and correlating treatment outcomes to its levels after therapy [20-25]. However, in a systematic review [11], it was shown that the presence or absence of Aa could not discriminate between subjects with aggressive periodontal disease from those with chronic periodontitis.

The present study focused on the presence of Aa as a determinant factor in some aspects of disease presentation. We observed a lack of association between Aa detection, as measured by HOMIM, and severity of clinical signs of the disease (i.e. pocket depth, clinical attachment levels at time of sampling). This is not unexpected, considering that the criteria for diagnosis of LAP used in this study are very strict for clinical signs of disease, although it is impossible to determine the disease state of the sampled sites, even before treatment is rendered. However, if we were to include healthy individuals or even healthy sites (i.e. $PD\leq 3mm$ and no bop), the presence of Aa could indeed be more correlated with the diseased clinical parameters [4,6,7,10,26]. It should be noted that the HOMIM assay has a sensitivity of about 0.1% and an overall limit of detection about 10^4 cells (Colombo, 2009). Consequently, it is possible that Aa at levels below the level of detection could still be present. Nevertheless, Aa-detected sites presented very high signal levels for (mean relative signal intensity was 3.81), which makes the two populations very distinct for Aa detection using HOMIM. Interestingly, at this detection level, the




Figure 2. Effect of Aa and microbial community composition, as measured by HOMIM hybridization scores. The following variables were chosen as descriptors of the subgingival biofilm community and were calculated for each sample: pocket depth (PD), and presence of Aa. This last term was chosen to determine if Aa alone reflects shifts in community composition. A. Non-metric multidimensional scaling analysis (NMDS) and B. Canonical correspondence analysis (CCA) ordination diagrams of microbial communities. Red circles indicate samples where Aa was detected by HOMIM, and black circles indicate samples where there was no Aa detected by HOMIM analysis. Aa was critical for determining bacterial community structures ($p < 0.01$).

doi: 10.1371/journal.pone.0085066.g002

presence or absence of Aa was associated with a difference in the composition of the microbial community as a whole.

We also evaluated the correlation between the detection of Aa and local inflammatory mediators as a measure of active disease. Interestingly, a higher expression of IL-8 in GCF was




Figure 3. Inflammatory mediator profile in the GCF, in both groups. Cytochemokines values are reported as normalized pg/mL. Box-plots show the mean (horizontal line), inter-quartile range (box), 5th -95th percentiles (vertical lines). *Statistically significant (p=0.01), t-test.

doi: 10.1371/journal.pone.0085066.g003

detected in *Aa*-non-detected sites as compared to *Aa*-detected sites. IL-8 is involved in the recruitment of PMNs and is highly expressed in the junctional epithelium adjacent under the conditions of health whereby disease has been shown to modulate these levels both positively and negatively. Many organisms, including *P.gingivalis*, have virulence factors which affect IL8 stability within the GCF. Our data indicate that the non-detection of *Aa* results in a different microbial community than when it is detected at levels above 10⁴, thus it is possible that the absence/minimal presence of *Aa*, i.e. *Aa*-non-detected sites promotes an environment for an organism that possess these virulence factors. As demonstrated in our results, *P.gingivalis* may be frequently detected when *Aa* is not, where

IL-8 levels were lower. Although *P.gingivalis* have been associated with periodontal inflammatory response, its potential to induce this response may shift according to the environment, specific virulent factors, and association with other species of the biofilm[27]. For instance, specific structures on *P.gingivalis* LPS may up-regulate or down-regulate levels of pro-inflammatory markers, including IL-8[28]. In addition, the presence of other bacterial species [29], as well as different peptides [30,31] may also reduce this organism's pro-inflammatory abilities. On the other hand, it is possible that *Aa* has a direct effect on the suppression of IL8 from the junctional epithelium. However, the differences in IL8 expression in the GCF did not correlate with differences in clinical parameters.




Figure 4. Systemic LPS-induced cyto/chemokine values in both groups. Values are reported as normalized log pg/mL. Box-plots show the mean (horizontal line), inter-quartile range (box), 5th -95th percentiles (vertical lines). *Indicates statistically significant differences between groups (Aa+ and Aa-) by t-test ($p < 0.05$). #Indicates statistically significant differences within groups regarding TLR stimulation by One-way ANOVA ($p < 0.05$).

doi: 10.1371/journal.pone.0085066.g004

Both *Aa-detected* and *Aa-non-detected* cohorts had already advanced clinical signs of LAP, thus IL8 levels may be more indicative of active vs. inactive disease than severity of disease, which would also suggest that the presence and/or absence of *Aa* may also be a correlate of disease activity, although data presented here are insufficient to support this

hypothesis. Longitudinal studies of the disease need to be conducted to assess this hypothesis.

Elevated local inflammation can sometimes result in the dissemination of bacteria and their products systemically. Interestingly, no differences in the levels of plasma LPS was observed between *Aa-detected* and *Aa-non-detected* cohorts. This was not unexpected considering that other than IL8, which

does not directly contribute to epithelial ulceration, there were no significant differences in the levels of cyto/chemokines within the GCF of the *Aa-detected* and *Aa-non-detected* cohorts.

We have previously published that LAP individuals present with a hyper-inflammatory phenotype whereby they respond in a more robust manner to stimuli [8]. Here we demonstrate that overall whether an individual is colonized with *Aa* or not does not correlate with their level of LPS-induced responsive potential, with one exception. As a group, individuals with diseased sites that have *Aa detected*, respond more robustly to *Ec* LPS-stimulation as measured by IL1 β and TNF α . *E.coli* LPS is mainly a TLR4 agonist, as is *Aa* LPS [32] [33] and this is why it was chosen for evaluation in the present study. Thus, one could hypothesize that in *Aa* carriers the potential for hyper-secretion of TNF α and IL1 β could result in more robust tissue damage and disease progression. However, we did not observe elevated levels of TNF α nor IL1 β in the GCF of *Aa detected* sites, nor was the snapshot of disease severity different in those individuals with *Aa-detected* or *Aa-non-detected* sites. The direct relationship between *Aa* presence, GCF markers and host inflammatory response to LPS cannot be resolved in this cross-sectional study but could be determined in future studies that test the likelihood that these cytokine elevations could be found prior to disease or during disease progression at the local site.

In summary, the present study demonstrates that, in this population, similar clinical parameters of disease were

observed regardless of whether *Aa* was detected or remained undetected by HOMIM in diseased sites. This study is cross-sectional and thus, it only represents a snapshot of the disease state at one point in time. However, our findings indicate potential differences in the microbial community profile associated (*G.elegans* and *P.loeschii*) or not associated (*P.gingivalis* and *P.micra*) with this species as well as effects on the local inflammation (IL8 in GCF). Interestingly, individuals with detectable levels of *Aa* also present with a TLR4 (*Ec* LPS) induced TNF α and IL1 β hyper-responsiveness. Further longitudinal studies are necessary to evaluate whether these differences could impact disease progression and treatment responses.

Acknowledgements

We thank Drs. Edward Zapert, John Bidwell and Dan Cober and all the staff from Leon County Dental Clinic, for the clinical assistance with this study. Clinical trial registration: #NCT01330719.

Author Contributions

Conceived and designed the experiments: LMS SMW BP IA. Performed the experiments: HH PFG. Analyzed the data: PFG HH VKC. Contributed reagents/materials/analysis tools: SMW VKC LMS. Wrote the manuscript: PFG LMS.

References

- Lamell CW, Griffen AL, McClellan DL, Leys EJ (2000) Acquisition and colonization stability of *Actinobacillus actinomycetemcomitans* and *Porphyromonas gingivalis* in children. *J Clin Microbiol* 38: 1196-1199. PubMed: 10699021.
- Moore WE, Moore LH, Ranney RR, Smibert RM, Burmeister JA et al. (1991) The microflora of periodontal sites showing active destructive progression. *J Clin Periodontol* 18: 729-739. doi:10.1111/j.1600-051X.1991.tb00064.x. PubMed: 1752997.
- Socransky SS, Haffajee AD (1992) The bacterial etiology of destructive periodontal disease: current concepts. *J Periodontol* 63: 322-331. doi: 10.1902/jop.1992.63.4s.322. PubMed: 1573546.
- Fine DH, Markowitz K, Furgang D, Fairlie K, Ferrandiz J et al. (2007) *Aggregatibacter actinomycetemcomitans* and its relationship to initiation of localized aggressive periodontitis: longitudinal cohort study of initially healthy adolescents. *J Clin Microbiol* 45: 3859-3869. doi: 10.1128/JCM.00653-07. PubMed: 17942658.
- Armitage GC (1999) Development of a classification system for periodontal diseases and conditions. *Ann Periodontol* 4: 1-6. doi: 10.1902/annals.1999.4.1.1. PubMed: 10863370.
- Haubek D (2010) The highly leukotoxic JP2 clone of *Aggregatibacter actinomycetemcomitans*: evolutionary aspects, epidemiology and etiological role in aggressive periodontitis. *APMIS Suppl*: 1-53. PubMed: 21214629.
- Haubek D, Ennibi OK, Poulsen K, Vaeth M, Poulsen S et al. (2008) Risk of aggressive periodontitis in adolescent carriers of the JP2 clone of *Aggregatibacter (Actinobacillus) actinomycetemcomitans* in Morocco: a prospective longitudinal cohort study. *Lancet* 371: 237-242. doi: 10.1016/S0140-6736(08)60135-X. PubMed: 18207019.
- Shaddox L, Wiedey J, Bimstein E, Magnuson I, Clare-Salzler M et al. (2010) Hyper-responsive phenotype in localized aggressive periodontitis. *J Dent Res* 89: 143-148. doi: 10.1177/0022034509353397. PubMed: 20042739.
- Shaddox LM, Wiedey J, Calderon NL, Magnuson I, Bimstein E et al. (2011) Local inflammatory markers and systemic endotoxin in aggressive periodontitis. *J Dent Res* 90: 1140-1144. doi: 10.1177/0022034511413928. PubMed: 21730256.
- Shaddox LM, Huang H, Lin T, Hou W, Harrison PL et al. (2012) Microbiological Characterization in Children with Aggressive Periodontitis. *J Dent Res* 91: 927-933. PubMed: 22863892.
- Mombelli A, Casagni F, Madianos PN (2002) Can presence or absence of periodontal pathogens distinguish between subjects with chronic and aggressive periodontitis? A systematic review. *J Clin Periodontol* 29 (Suppl 3): 10-37; discussion:
- Lang NP, Bartold M, Cullinan M, Jeffcoat M, Mombelli A et al. (1999) Consensus Report: Aggressive Periodontitis. 53 p.
- Colombo AP, Boches SK, Cotton SL, Goodson JM, Kent R et al. (2009) Comparisons of subgingival microbial profiles of refractory periodontitis, severe periodontitis, and periodontal health using the human oral microbe identification microarray. *J Periodontol* 80: 1421-1432. doi: 10.1902/jop.2009.090185. PubMed: 19722792.
- Rothman KJ (1990) No adjustments are needed for multiple comparisons. *Epidemiology* 1: 43-46. doi: 10.1097/00001648-199001000-00010. PubMed: 2081237.
- Hornik K, Leisch F, Team RDC (2005) R version 2.1.0 (vol 19, pg 661, 2004). *Computational Statistics* 20: 197-202. doi:10.1007/BF02789699.
- Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate - A practical and powerful approach to multiple testing *Journal of the Royal Statistical Society Series B-Methodological* 57.
- Gotelli NJ, Ellison AM (2004). *A primer of ecological statistics*.
- Sanders NJ, Gotelli NJ, Heller NE, Gordon DM (2003) Community disassembly by an invasive species. *Proc Natl Acad Sci U S A* 100: 2474-2477. PubMed: 12604772.
- Klepac-Ceraj V, Lemon KP, Martin TR, Allgaier M, Kembel SW et al. (2010) Relationship between cystic fibrosis respiratory tract bacterial communities and age, genotype, antibiotics and *Pseudomonas aeruginosa*. *Environ Microbiol* 12: 1293-1303. doi:10.1111/j.1462-2920.2010.02173.x. PubMed: 20192960.
- Haffajee AD, Socransky SS, Ebersole JL, Smith DJ (1984) Clinical, microbiological and immunological features associated with the treatment of active periodontosis lesions. *J Clin Periodontol* 11: 600-618. doi:10.1111/j.1600-051X.1984.tb00913.x. PubMed: 6386896.
- Christersson LA, Slots J, Rosling BG, Genco RJ (1985) Microbiological and clinical effects of surgical treatment of localized juvenile

- periodontitis. *J Clin Periodontol* 12: 465-476. doi:10.1111/j.1600-051X.1985.tb01382.x. PubMed: 3894435.
22. Mandell RL, Tripodi LS, Savitt E, Goodson JM, Socransky SS (1986) The effect of treatment on *Actinobacillus actinomycetemcomitans* in localized juvenile periodontitis. *J Periodontol* 57: 94-99. doi:10.1902/jop.1986.57.2.94. PubMed: 2420958.
 23. Kaner D, Christan C, Dietrich T, Bernimoulin JP, Kleber BM et al. (2007) Timing affects the clinical outcome of adjunctive systemic antibiotic therapy for generalized aggressive periodontitis. *J Periodontol* 78: 1201-1208. doi:10.1902/jop.2007.060437. PubMed: 17608574.
 24. Leung WK, Ngai VK, Yau JY, Cheung BP, Tsang PW et al. (2005) Characterization of *Actinobacillus actinomycetemcomitans* isolated from young Chinese aggressive periodontitis patients. *J Periodontol Res* 40: 258-268. doi:10.1111/j.1600-0765.2005.00805.x. PubMed: 15853973.
 25. Zambon JJ, Christersson LA, Slots J (1983) *Actinobacillus actinomycetemcomitans* in human periodontal disease. Prevalence in patient groups and distribution of biotypes and serotypes within families. *J Periodontol* 54: 707-711. doi:10.1902/jop.1983.54.12.707. PubMed: 6358452.
 26. Ennibi OK, Benrachadi L, Bouziane A, Haubek D, Poulsen K (2012) The highly leukotoxic JP2 clone of *Aggregatibacter actinomycetemcomitans* in localized and generalized forms of aggressive periodontitis. *Acta Odontol Scand* 70: 318-322. doi:10.3109/00016357.2011.642002. PubMed: 22251014.
 27. Cugini C, Klepac-Ceraj V, Rackaityte E, Riggs JE, Davey ME (2013) *Porphyromonas gingivalis*: keeping the pathos out of the biont. *J Oral Microbiol* 5. PubMed: 23565326.
 28. Herath TD, Darveau RP, Seneviratne CJ, Wang CY, Wang Y et al. (2013) Tetra- and penta-acylated lipid A structures of *Porphyromonas gingivalis* LPS differentially activate TLR4-mediated NF-kappaB signal transduction cascade and immuno-inflammatory response in human gingival fibroblasts. *PLOS ONE* 8: e58496. doi:10.1371/journal.pone.0058496. PubMed: 23554896.
 29. Zhao JJ, Feng XP, Zhang XL, Le KY (2012) Effect of *Porphyromonas gingivalis* and *Lactobacillus acidophilus* on secretion of IL1B, IL6, and IL8 by gingival epithelial cells. *Inflammation* 35: 1330-1337. doi:10.1007/s10753-012-9446-5. PubMed: 22382516.
 30. Kraus D, Winter J, Jepsen S, Jäger A, Meyer R et al. (2012) Interactions of adiponectin and lipopolysaccharide from *Porphyromonas gingivalis* on human oral epithelial cells. *PLOS ONE* 7: e30716. doi:10.1371/journal.pone.0030716. PubMed: 22319581.
 31. Inomata M, Into T, Murakami Y (2010) Suppressive effect of the antimicrobial peptide LL-37 on expression of IL-6, IL-8 and CXCL10 induced by *Porphyromonas gingivalis* cells and extracts in human gingival fibroblasts. *Eur J Oral Sci* 118: 574-581. doi:10.1111/j.1600-0722.2010.00775.x. PubMed: 21083618.
 32. Mahanonda R, Pichyangkul S (2007) Toll-like receptors and their role in periodontal health and disease. *Periodontol* 2000 43: 41-55. doi:10.1111/j.1600-0757.2006.00179.x. PubMed: 17214834.
 33. Lee SH, Kim KK, Rhyu IC, Koh S, Lee DS et al. (2006) Phenol/water extract of *Treponema socranskii* subsp. *socranskii* as an antagonist of Toll-like receptor 4 signalling. *Microbiology* 152: 535-546. doi:10.1099/mic.0.28470-0. PubMed: 16436441.